

NUCLEIC ACID ENZYME FOR RNA CLEAVAGE

This application is a continuation of
PCT/CA99/00391, filed April 29, 1999, which claims priority to
Canadian patent application 2,230,203, filed April 29, 1998,
5 both of which are incorporated herein in their entirety.

TECHNICAL FIELD

The invention relates to a novel ribozyme
construction for the specific recognition and cleavage of RNA,
and biotechnological as well as therapeutic uses thereof.

BACKGROUND ART

Though enzymatic activity has long been considered
the exclusive domain of proteins, discoveries in molecular
biology over the past couple of decades have led to the
realization that ribonucleic acid (RNA) can also function as
15 an enzyme. RNA enzymes are often referred to as ribozymes.

Ribozyme substrates are generally confined to RNA
molecules, and enzymatic activities of ribozymes include the
cleavage and/or ligation of RNA molecules. The cleavage
activity may be intramolecular, known as *cis*-acting or
20 intermolecular, known as *trans*-acting. There are at least
five classes of ribozymes known, including Group I introns,
Group II introns, hammerhead, hairpin, and delta ribozymes.
The last three are derived from plant satellites and viroids.

Since 1982, several unexpected diseases caused by
25 RNA-based pathogenic agents have emerged. These include the
lethal Acquired Immune Deficiency Syndrome (AIDS) and delta
hepatitis, a particularly virulent form of fulminant hepatitis
caused by a viroid-like RNA agent. These blood-borne diseases
are spread at the RNA level, manifest themselves in cells of
30 patients, and are by now present within the bloodstream of
millions of individuals. Conventional biotechnology, with its
reliance on recombinant DNA methods and DNA-level intervention
schemes, has been slow to provide valid approaches to combat
these diseases.

0969667 "103000

Two forms of delta ribozymes, namely genomic and antigenomic, are derived, and referred to by, the polarity of the hepatitis delta virus (HDV) genome from which the ribozyme is generated. Like hammerhead and hairpin ribozymes, the delta ribozymes cleave a phosphodiester bond of their RNA substrates and give rise to reaction products containing a 5'-hydroxyl and a 2',3'-cyclic phosphate termini. They are metalloenzymes and a low concentration (<1 mM) of magnesium (Mg^{2+}) or calcium (Ca^{2+}) is required for delta ribozyme cleavage. Both genomic strand and antigenomic strand forms exhibit self-cleavage activity, and it has been suggested that they are involved in the process of viral replication (Lazinski, D. W., and Taylor, J. M. (1995) *RNA* 1, 225-233).

Delta ribozymes derived from the genome of HDV are of interest in the development of a gene regulation system in which the designed ribozymes would down-regulate the expression of a target gene. The facts that delta ribozymes are derived from HDV and that this pathogen naturally replicates in animal systems, suggest that this catalytic RNA could be used to control gene expression in human cells. Like other ribozymes, the designed ribozyme should specifically cleave its target substrates while leaving other cellular RNA molecules intact.

Trans-acting ribozymes carry out intermolecular cleavage activity. Some trans-acting delta ribozymes have been developed by removing a single-stranded junction which connects the catalytic portion to the substrate portion in cis-acting delta ribozymes. This results in two separate molecules, one possessing the substrate sequence and the other the catalytic property (Been, M.D. and Wichhan, G.S. (1997) *Eur. J. Biochem.*, 247, 741-753). Interactions between such delta ribozymes and the substrate occur through the formation of a helix, referred as the P1 stem. However, the example of the trans-acting ribozyme disclosed by Been et al. (supra) was not useful for cleaving long substrate molecules, such as those having therapeutic applications.

In United States Patent No. 5,225,337, issued on

09699667 103000

conditions which involve RNA expression, such as AIDS. These ribozymes consist of at least 18 consecutive nucleotides from the conserved region of HDV isolates between residues 611 and 771 on the genomic strand and between residues 845 and 980 on the complementary antigenomic strand. These ribozymes are proposed to fold into an axe-head model secondary structure (Branch, A. D., and Robertson, H. D. (1991) *Proc. Natl .Acad. Sci. USA* **88**, 10163-10167). The ribozymes developed according to this model structure require the substrate to be bound to the ribozyme through the formation of two helices, one located on either side of the cleavage site. Further, such ribozymes apparently require a 12-15 nucleotide recognition sequence in the substrate in order to exhibit the desired activity. Such a long recognition sequence is not practical in the development of therapeutic or diagnostic applications.

In United States Patent No. 5,625,047, issued on April 29, 1997 in the names of Michael D. Been et al., there are disclosed enzymatic RNA molecules proposed to fold into a pseudoknot model secondary structure (discussed below). The method disclosed for the development of efficient ribozymes requires a short recognition sequence of only 7 to 8 nucleotides in the substrate, a preference for a guanosine base immediately 3' to the cleavage site, a preference for U, C or A immediately 5' to the cleavage site, and the availability of a 2'-hydroxyl group for cleavage to occur. Thus, the specificity of recognition of these ribozymes is limited to 6 or 7 base pairing nucleotides with the substrate and a preference of the first nucleotide located 5' to the cleavage site. Neither tertiary interaction(s) between the base paired nucleotides and another region of the ribozyme, nor single-stranded nucleotides are involved to define the specificity of recognition of these ribozymes. Because the recognition features are limited, these ribozymes have a

limited specificity, and thus, are not practical for further clinical or biotechnical applications.

A pseudoknot-like structure for *delta* ribozymes has been proposed by Perrotta and Been (Perrotta, A. T., and Been, M. D. (1991) *Nature* 350, 434-436). This model structure consists of two stems (P1 and P2), two stem-loops (P3 and P4) and three single-stranded regions (J1/2, J1/4 and J4/2). An additional stem, named P1.1, has been formed by two GC base pairs between nucleotides from the J1/4 junction and the P3 loop (Ferré-D'Amaré, A.R., Zhou, K. and Doudna, J.A. (1998) *Nature*, 350, 434-436).

It would be highly desirable to be provided with a novel *delta* ribozyme for the cleavage of both small and large RNA substrates for which the specificity of recognition is well defined. Such specificity would yield optimal conditions for further therapeutical and biotechnological developments of *delta* ribozymes.

SUMMARY OF THE INVENTION

One aim of the present invention is to provide a novel *delta* ribozyme for the cleavage of RNA substrates for which the specificity is defined by a domain composed of at least 7 nucleotides. It is also an aim to provide a method for the development of such ribozymes.

In one aspect, the invention provides a method for cleaving a nucleic acid substrate with a nucleic acid enzyme at a cleavage site comprising mixing the substrate with the enzyme, wherein the substrate includes a 7 nucleotide sequence with at least 6 nucleotides 3' to the cleavage site and at least 1 nucleotide 5' to the cleavage site of formula:



wherein each

N is a nucleotide which may be the same or different,
H is a nucleotide selected from the group consisting of A, U, C, and T, and

is the site of cleavage, and
H' is a ribonucleotide selected from the group consisting
of A, U, and C,

wherein

5 (i) the first nucleotide 3' to the cleavage site is
capable of forming a wobble pair with the enzyme,

(ii) the second, third, fifth, and sixth nucleotides 3'
to the cleavage site are capable of forming conventional
Watson-Crick base pairs with the enzyme,

10 (iii) the fourth nucleotide 3' to the cleavage site is
capable of forming a triplet with the enzyme comprising a non-
conventional Watson-Crick base pair and a conventional Watson-
Crick base pair, and

(iv) the ribonucleotide directly 5' to the cleavage site
15 does not form a base pair with the enzyme; and
the enzyme comprises a substrate binding portion which is
capable of base pairing to the 6 nucleotides 3' to the
cleavage site of the substrate and which binding portion
comprises the sequence:

20

3'-UNNXNN-5'

wherein each

25 N is a nucleotide which may be the same or different, and
X is a nucleotide selected from the group consisting of
T, U, A, and G,

whereby binding of the substrate to the enzyme effects
cleavage of the substrate at the cleavage site.

30 In another aspect, the invention provides a nucleic
acid enzyme capable of recognizing and cleaving a nucleic acid
substrate at a cleavage site comprising a substrate binding
portion which is capable of base pairing to the 6 nucleotides
3' to the cleavage site of the substrate and which binding
portion comprises the sequence:


35

3'-UNNXNN-5'

09699667 "103000"

N is a nucleotide which may be the same or different, and
X is a nucleotide selected from the group consisting of
T, U, A, G, and

10



15

is the site of cleavage, and
H' is a ribonucleotide selected from the group consisting
of A, U, and C,

wherein

(ii) the second, third, fifth, and sixth nucleotides 3' to the cleavage site are capable of forming conventional Watson-Crick base pairs with the enzyme,

(iv) the first ribonucleotide directly 5' to the cleavage site does not form a base pair with the enzyme.

30 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the secondary structure and nucleotide sequences of two trans-acting antigenomic delta ribozymes of the invention and complementary substrates; panel A is the secondary structure of the complex formed between δ RzPl.1 and a substrate Sp1.1; panel B is the P1 region of the

Figure 2 illustrates the secondary structure of a ribozyme in accordance with the invention, with an ultrastable L4 loop; in the inset is the sequence of a 14-nucleotide long substrate;

10 Figure 4 shows a two-dimensional representation of a
catalytic trimolecular complex (RzA: RzB:S) of the invention;

Figure 6 shows a two-dimensional representation of a catalytic trimolecular complex (RzA: RzB:S); the influence of 2'-OH groups individually at positions 9 to 15 on RzB by replacing the ribonucleotide at these positions with the corresponding deoxy-ribonucleotide is shown; the symbol - represents a two-fold diminution of activity compared to an unmodified RzB while the symbol = represents an unchanged catalytic activity; symbols + and ++ respectively represent an increased activity of 1.5- and 2- fold; horizontal bars represent base pairs; wobble and homopurine base pairs are respectively represented by one and two ovals; the arrow indicates the site of catalytic cleavage;

30 (AAA) species found at the 5' ends of the minus and plus DNA strands, respectively; the dashed line indicates the presence of the single stranded gap; the RNA products are depicted by wavy lines; the target area is located in pre-S2 and S regions, and is indicated by the scissors symbol; panel B
35 illustrates the secondary structure of an engineered ribozyme of the the invention, such that the substrate binding region

[illegible]

5

10

15

25

30

35

sequence with at least 6 nucleotides 3' to the cleavage site and at least 1 nucleotide 5' to the cleavage site of formula:



wherein each

N is a nucleotide which may be the same or different,
H is a nucleotide selected from the group consisting of
A, U, C, and T, and

10 is the site of cleavage, and

H' is a ribonucleotide selected from the group consisting of A, U, and C.

The first nucleotide 3' to the cleavage site is capable of forming a wobble pair with the enzyme. The wobble base pair (G-U) at the cleavage site is required to maintain a high level of cleavage. Conventional Watson-Crick base pairs such as A-U and G-C, as well as mismatches at this position decrease the cleavage activity.

The second, third, fifth, and sixth nucleotides 3' to the cleavage site are capable of forming conventional Watson-Crick base pairs with the enzyme.

The fourth nucleotide 3' to the cleavage site is capable of forming a conventional Watson-Crick base pair with the substrate binding region of the enzyme. Additionally, such base pair interacts with a nucleotide elsewhere in the ribozyme (i.e. the nucleic acid enzyme) to form a triplet by means of a non-conventional Watson-Crick base pair. Non-conventional Watson-Crick base pairs include Hoogsteen pairs and reversed-Hoogsteen pairs. The position requires an A, U, or C.

The ribonucleotide directly 5' to the cleavage site does not form a base pair with the ribozyme.

Preferably, the substrate molecule does not contain two consecutive pyrimidine nucleotides directly 5' to the cleavage site.

In another preferred aspect, the substrate comprises the sequence 5'-H' GNNHNNN-3', more preferably the sequence 5'-NNRH' GNNHNNN-3', wherein R is G or A.

In one embodiment, the substrate preferably
5 comprises the sequence 5'-RRRH' GNNHNNN-3'. More preferably, such sequence is selected from the group consisting of 5'-GGGC GNNUNNN-3', 5'-GGGC GNNCNNN-3', 5'-GGGU GNNUNNN-3', 5'-GGGU GNNCNNN-3', and 5'-AAAC GNNUNNN-3'.

In another embodiment, the substrate preferably
10 comprises the sequence 5'-YHRH' GNNHNNN-3', wherein Y is C, U, or T. It is preferred that the four nucleotides directly 5' to the cleavage site are chosen such that Y is C or U, preferably C; H is one of U, C, or A, preferably U or C, more preferably U; R is preferably A; and H is A, C, or U,
15 preferably A or C, more preferably A.

It is preferable that the four nucleotides directly 5' to the cleavage site do not form a hairpin structure.

Selection of Ribozyme Sequence

By ribozymes, it is meant a nucleic acid enzyme, in
20 other words any nucleic acid sequence having enzymatic activity, i.e. the ability to catalyze a reaction. As such it includes nucleic acid sequences made up of ribonucleotides, deoxyribonucleotides, or mixtures of both. Nucleotides may also include synthetic or modified nucleotides.

25 The selection of the sequence of the substrate binding region of the ribozyme, should be done such that the binding region comprises the sequence 3'-UNNXNN-5', wherein each N is a nucleotide which may be the same or different, and X is a nucleotide selected from the group consisting of T, U,
30 A, and G.

The invention preferably provides for a nucleic acid enzyme with a secondary structure which comprises three or more distinct double-stranded regions, or stem-regions. This includes regions of base-pairing which may or may not be
35 capped by a single-stranded loop, to form a stem-loop region. Preferably, the nucleic acid ribozyme includes two or more distinct single-stranded regions, one of which includes a

05699667 "103000

The invention preferably contemplates the use of
5 nucleic acid enzymes derived from hepatitis delta virus, known
as *delta* ribozymes.

Trans-acting delta ribozymes of the invention were generated based on the pseudoknot-like structure proposed by Perrotta and Been, by removing the single-stranded region (region J1/2) located at the junction between the P1 and P2 stems. In addition, the P2 stem was elongated, by introducing, for instance, three G-C base pairs, and by shortening the P4 stem.

In another aspect, the invention provides for a
30 ribozyme with an elongated P2 stem and shortened P4 stem,
which further comprises a modification of the L4 loop.
Figures 2 and 3 show ribozymes in accordance with this
embodiment. S and Rz represent substrate and ribozyme
respectively.

35 In one aspect, the invention provides for a
bimolecular ribozyme. This may be achieved by removal of the

L4 loop. Figure 4 shows a ribozyme in accordance with this embodiment.

Applications

Ribozyme activity can be optimized by chemically synthesizing ribozymes with modifications that prevent their degradation by serum ribonucleases (see e.g., Eckstein et al., International Publication No. WO 92/07065; Perreault et al., Nature 1990, 344:565; Pieken et al., Science 1991, 253:314; and Chowrira et al., 1993 J. Biol. Chem. 268, 19458, which describe various chemical modifications that can be made to the sugar moieties of enzymatic RNA molecules, all of which publications are incorporated by reference herein), modifications which enhance their efficacy in cells, and removal of helix-containing bases to shorten RNA synthesis times and reduce chemical requirements.

In one aspect, the invention provides a substrate molecule which is a target RNA, such as a viral RNA, or an RNA crucial to the life cycle of a pathogen, or an RNA manifested as a result of an inherited disease, based on the substrate specificity described herein.

Ribozymes are added directly, or can be complexed with cationic lipids, packaged within liposomes, or otherwise delivered to target cells. The RNA or RNA complexes can be locally administered to relevant tissues *ex vivo*, or *in vivo* through injection, aerosol inhalation, infusion pump or stent, with or without their incorporation in biopolymers.

Sullivan, et al., (WO 94/02595, incorporated by reference herein), describes general methods for delivery of enzymatic RNA molecules. Ribozymes may be administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres. For some indications, ribozymes may be directly delivered *ex vivo* to cells or tissues with or without the aforementioned vehicles. Alternatively, the RNA/vehicle combination is locally

000007 29966660

delivered by direct injection or by use of a catheter,
infusion pump or stent. Other routes of delivery include, but
are not limited to, intravascular, intramuscular, subcutaneous
or joint injection, aerosol inhalation, oral (tablet or pill
5 form), topical, systemic, ocular, intraperitoneal and/or
intrathecal delivery. More detailed descriptions of ribozyme
delivery and administration are provided in Sullivan, et al.,
("Method and Reagent for Treatment of Arthritic Conditions"
U.S.S.N. 08/152,487, filed November 12, 1993, and incorporated
10 by reference herein).

Another means of accumulating high concentrations of
a ribozyme(s) within cells is to incorporate the ribozyme-
encoding sequences into a DNA expression vector. Transcription
of the ribozyme sequences are driven from a promoter for
15 eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol
II), or RNA polymerase III (pol III). Transcripts from pol II
or pol III promoters will be expressed at high levels in all
cells; the levels of a given pol II promoter in a given cell
type will depend on the nature of the gene regulatory
20 sequences (enhancers, silencers, etc.) present nearby.
Prokaryotic RNA polymerase promoters are also used, providing
that the prokaryotic RNA polymerase enzyme is expressed in the
appropriate cells (Elroy-Stein, O. and Moss, B., 1990, Proc.
Natl. Acad. Sci. U S A, 87, 6743-7; Gao, X. and Huang, L.,
25 1993, Nucleic Acids Res., 21, 2867-72; hereby incorporated by
reference). Several investigators have demonstrated that
ribozymes expressed from such promoters can function in
mammalian cells (e.g. Kashani-Sabet, M., et al., 1992,
Antisense Res. Dev., 2, 3-15; Ojwang, J. O., et al., 1992,
30 Proc. Natl. Acad. Sci. U S A, 89, 10802-6; hereby incorporated
by reference). The above ribozyme transcription units can be
incorporated into a variety of vectors for introduction into
mammalian cells, including but not restricted to, plasmid DNA
vectors, viral DNA vectors (such as adenovirus or
35 adeno-associated vectors), or viral RNA vectors (such as
retroviral, Semliki forest virus, hepatitis delta virus, and
sindbis virus vectors).

0969667-103000

Thus, ribozymes of the present invention that cleave target mRNA and thereby inhibit and/or reduce target activity have many potential therapeutic uses, and there are reasonable modes of delivering the ribozymes in a number of the possible indications.

By "inhibit" is meant that the activity or level of target RNA is reduced below that observed in the absence of the ribozyme, and preferably is below that level observed in the presence of an inactive RNA molecule able to bind to the same site on the RNA, but unable to cleave that RNA.

By "vectors" is meant any nucleic acid and/or viral-based construct used to deliver a desired nucleic acid.

Examples

Example 1: Preparation of Ribozymes, Substrates, and Plasmids.

Construction of plasmids carrying ribozymes of the invention. The antigenomic ribozyme sequence of the hepatitis delta virus described by Makino et al (Makino, S. et al. (1987) *Nature* 329, 343-346, hereby incorporated by reference) was used as the basis for generating trans-acting delta ribozymes of the invention. Briefly, the construction was performed as follows. Two pairs of complementary and overlapping oligonucleotides, representing the entire length of the ribozyme (57 nt), were synthesized and subjected to an annealing process prior to cloning into pUC19. The annealed oligonucleotides were ligated to *Hind*III and *Sma*I co-digested pUC19 to give rise to a plasmid harboring the delta ribozyme (referred to as p δ RzP1.1). The minigene was designed so as to have unique *Sph*I and *Sma*I restriction sites. The sequence of the T7 RNA promoter was included at the 5' end of the ribozyme so as to permit *in vitro* transcription. Variations based on this "wild type" ribozyme are constructed by replacing the *Sph*I-*Sma*I fragment of p δ RzP1.1 by an oligonucleotide duplex containing the desired sequence. The sequences of engineered ribozymes were confirmed by DNA sequencing. Plasmids contain-

09699667-103000

ing wild type and mutant ribozymes were then prepared using Qiagen tip-100 (Qiagen Inc.), digested with *Sma*I, purified by phenol and chloroform extraction and precipitated for further use as templates for in vitro transcription reactions.

5 *Synthesis of Ribozymes and Substrates. Ribozyme:*

In vitro transcription reactions contained 5 µg linearized recombinant plasmid DNA as template, 27 units RNAGuard (RNase inhibitor (Pharmacia), 4 mM of each rNTP (Pharmacia), 80 mM HEPES-KOH pH 7.5, 24 mM MgCl₂, 2 mM spermidine, 40 mM DTT, 0.01 unit Pyrophosphatase (Boehringer Mannheim) and 25 µg purified T7 RNA polymerase in a final volume of 50 µL, and were incubated at 37°C for 4 hr. *Substrates:* Deoxyoligonucleotides (500 pmoles) containing the substrate and the T7 promoter sequence were denatured by heating at 95°C for 5 min in a 20 µL mixture containing 10 mM Tris-HCl pH 7.5, 10 mM MgCl₂, 50 mM KCl₂, and allowed to cool slowly to 37°C. The *in vitro* transcription reactions were carried out using the resulting partial duplex formed as template under the same conditions as described for the production of the ribozyme.

20 After incubation, the reaction mixtures were fractionated by denaturing 20% polyacrylamide gel electrophoresis (PAGE, 19:1 ratio of acrylamide to bisacrylamide) containing 45 mM Tris-borate pH 7.5, 7 M urea and 1 mM EDTA. The reaction products were visualized by UV shadowing. The bands corresponding to the correct sizes of either ribozymes or substrates were cut out, and the transcripts eluted overnight at 4°C in a solution containing 0.1% SDS and 0.5 M ammonium acetate. The transcripts were then precipitated by the addition of 0.1 vol 3 M sodium acetate pH 5.2 and 2.2 vol ethanol. Transcript yield was determined by spectrophotometry.

35 *Synthesis and Purification of RNA and RNA/DNA Mixed Polymer:* RNA and RNA-DNA mixed polymers were synthesized on an automated oligonucleotide synthesizer, and deprotected according to previously described procedures (Perreault, J.P., and Altman, S. (1992) J. Mol. Biol. 226, 339-409 hereby incorpor-

09699667 103000

ated by reference). These polymers were purified by 20% PAGE. Major bands were excised and eluted as described above.

End-labelling of RNA with [γ - 32 P]ATP. Purified transcripts (10 pmoles) were dephosphorylated in a 20 μ L reaction mixture containing 200 mM Tris-HCl pH 8.0, 10 units RNA guard, and 0.2 unit calf intestine alkaline phosphatase (Pharmacia). The mixture was incubated at 37°C for 30 min, and then extracted twice with a same volume of phenol:chloroform (1:1). Dephosphorylated transcripts (1 pmole) were end-labelled in a mixture containing 1.6 pmole [γ - 32 P]ATP, 10 mM Tris-HCl pH 7.5, 10 mM MgCl₂, 50 mM KCl and 3 units T4 polynucleotide kinase (Pharmacia) at 37°C for 30 min. Excess [γ - 32 P]ATP was removed by applying the reaction mixture onto a spin column packed with a G-50 Sephadex gel matrix (Pharmacia). The concentration of labelled transcripts was adjusted to 0.01 pmol per mL by the addition of water.

Example 2: Kinetics

Cleavage reactions. To initiate a cleavage reaction, various concentrations of ribozymes were mixed with trace amounts of substrate (final concentration <1 nM) in a 18 μ L reaction mixture containing 50 mM Tris-HCl pH 7.5, and subjected to denaturation by heating at 95°C for 2 min. The mixtures were quickly placed on ice for 2 min and equilibrated to 37°C for 5 min prior to the initiation of the reaction. Unless stated otherwise, cleavage was initiated by the addition of MgCl₂ to 10 mM final concentration. The cleavage reactions were incubated at 37°C, and followed for 3.5 hours or until the endpoint of cleavage was reached. The reaction mixtures were periodically sampled (2-3 μ L), and these samples were quenched by the addition of 5 μ L stop solution containing 95% formamide, 10 mM EDTA, 0.05% bromophenol blue and 0.05% xylene cyanol. The resulting samples were analyzed by a 20% PAGE as described above. Both the substrate and the reaction product bands were detected using a Molecular Dynamic

radioanalytic scanner after exposition of the gels to a phosphoimaging screen.

Kinetic analysis. Measurement of pseudo-first-order rate constant (k_{cat} , K_M and k_{cat}/K_M) were performed under single turnover conditions. Briefly, trace amounts of end-labelled substrate (<1 nM) were cleaved by various ribozyme concentrations (5 to 500 nM). The fraction cleaved was determined, and the rate of cleavage (k_{obs}) obtained from fitting the data to the equation $A_t = A_{\infty}(1 - e^{-kt})$ where A_t is the percentage of cleavage at time t , A_{∞} is the maximum percent cleavage (or the end point of cleavage), and k is the rate constant (k_{obs}). Each rate constant was calculated from at least two measurements. The values of k_{obs} obtained were then plotted as a function of ribozyme concentrations for determination of the other kinetic parameters: k_{cat} , K_M and k_{cat}/K_M . Values obtained from independent experiments varied less than 15%. The requirement for Mg^{2+} by both ribozymes was studied by incubating the reaction mixtures with various concentrations of MgCl_2 (1 to 500 mM) in the presence of an excess of ribozyme (500 nM) over substrate (< 1nM). The concentrations of Mg^{2+} at the half maximal velocity were determined for both ribozymes. Determination of equilibrium dissociation constants (K_d). For mismatched substrates which could not be cleaved by the ribozyme, the equilibrium dissociation constants were determined. Eleven different ribozyme concentrations, ranging from 5 to 600 nM, were individually mixed with trace amounts of end-labelled substrates (< 1nM) in a 9 μL solution containing 50 mM Tris-HCl pH 7.5, heated at 95°C for 2 min and cooled to 37°C for 5 min prior to the addition of MgCl_2 to a final concentration of 10 mM, in a manner similar to that of a regular cleavage reaction. The samples were incubated at 37°C for 1.5 h, at which 2 μL of sample loading solution (50% glycerol, 0.025% of each bromophenol blue and xylene cyanol) was added, and the resulting mixtures were electrophoresed through a nondenaturing polyacrylamide gel (20% acrylamide with a 19:1

5 Quantification of bound and free substrates was performed following an exposure of the gels to a phosphoimaging screen as described earlier.

10 Specificity

15 ribozymes and substrates led to the characterizations of the
method for selecting the ribozyme and substrate sequences. A
summary of the kinetic data is given below.

20 the sequence 3'-UNNXNN-5'.

(11 nt) SP1.1 and SP1.2 for the kinetic studies (see Table 1).

25 in Fig. 1. $\delta\text{RzPl.2}$ differs from $\delta\text{RzPl.1}$ in that $\delta\text{RzPl.2}$ has two nucleotides, at positions 22 and 24 of $\delta\text{RzPl.1}$, interchanged (5'-CCCAGCU-3').

TABLE 1

Kinetic parameters	$\delta\text{RzP.1}$	$\delta\text{RzP.2}$
$k_{\text{cat}} (\text{min}^{-1})$	0.34 ± 0.02	0.13 ± 0.01
$K_{\text{M}}' (\text{nM})$	17.9 ± 5.6	16.7 ± 6.4
$k_{\text{cat}}/K_{\text{M}}' (\text{min}^{-1} \bullet \text{M}^{-1})$	1.89×10^7	0.81×10^7
$K_{\text{Mg}} (\text{mM})$	2.2 ± 1.0	2.1 ± 0.8

Table 1. Kinetic parameters of wild type ribozyme (δ RzP1.1) and mutant ribozyme (δ RzP1.2). Under single turnover conditions, trace amounts of end-labelled substrate (<1 nM) were cleaved by various concentrations of ribozyme (5 to 600 nM). Reactions carried out under these conditions displayed monophasic kinetics. The values were calculated from at least two independent experiments, and standard variations were less than 15%.

- 5
- 10 In order to compare the specificity of the delta ribozyme with various substrates, δ RzP1.1 was used under single turnover conditions as described above. The cleavage reactions were performed with a trace amount of each substrate (<1 nM) and 500 nM δ RzP1.1. Under these conditions, the
- 15 observed rates reflect the rates of cleavage without interference from either product dissociation or inhibition. For each substrate both the observed cleavage rate constants (k_{obs}) and the extent of cleavage were calculated and compared to those of the wild type substrate, as shown in Table 2.

09699667-1030000

TABLE 2

Table 2. Cleavage activity of shorter or mismatched substrates as compared to the wild type substrate (SP.1). Bold letters represent the nucleotides of wild type substrate recognized by δ RzP1.1. The numbers in subscript indicate the nucleotides of wild type substrate which were individually altered to generate shorter or mismatched substrates.

Substrates	Sequence	k_{obs}^a (min ⁻¹)	Extent of cleavage ^c (%)	k_{rel}^d	$\Delta\Delta G^{\text{Te}}$ (kcal/mol)
Wild type substrate (S11-mer)	GGGCG ₅ G ₆ G ₇ U ₈ C ₉ G ₁₀ G ₁₁	0.34 ± 0.02		1	-
S10-mer	GGGCGGGUCG	0.022 ± 0.01	28.8 ± 4.3	0.063	-1.69
S9-mer	GGGCGGGUC	na ^b	na ^b	-	-
S8-mer	GGGCGGGU	na ^b	na ^b	-	-
SG5A	GGGCGAGGUCGG	0.009 ± 0.002	20.0 ± 2.4	0.026	-2.25
SG5C	GGGCGGGUCGG	0.047 ± 0.017	1.7 ± 0.2	0.138	-1.22
SG6A	GGGCGAGGUCGG	0.026 ± 0.006	5.8 ± 0.5	0.076	-1.59
SG6U	GGGCGGUGUCGG	0.071 ± 0.026	3.7 ± 0.3	0.209	-0.96
SG7A	GGGCGGAUCGG	na ^b	na ^b	-	-
SG7U	GGGCGGUUCGG	na ^b	na ^b	-	-
SU8C	GGGCGGGCCGG	na ^b	na ^b	-	-
SU8G	GGGCGGGCCGG	na ^b	na ^b	-	-
SC9A	GGGCGGGUAGG	0.016 ± 0.007	8.2 ± 3.0	0.047	-1.88
SC9U	GGGCGGGUUGG	0.031 ± 0.005	21.2 ± 1.0	0.091	-1.48
SG10U	GGGCGGGUCUG	0.016 ± 0.002	8.4 ± 0.5	0.047	-1.88
SG11U	GGGCGGGUCGU	0.011 ± 0.001	32.1 ± 2.5	0.032	-2.12

^ak_{obs} is the observed rate of cleavage calculated from at least two measurements. ^bna represents no detectable cleavage activity after 3.5 hours incubation. ^cCleavage extent (%) is obtained by fitting the data to the equation $A_t = A_{\infty} (1 - e^{-kt})$, where A_t is the percentage of cleavage at time t , A_{∞} is the maximum percentage of the cleavage, and k is the rate constant. ^dk_{rel} is the relative rate constant as compared to that of wild type substrate. ^eΔG[‡], the apparent free energy of transition-state stabilization, was calculated using the equation $\Delta\Delta G^{\ddagger} = RT \ln k_{rel}$, where $T = 310.15 \text{ K (37}^{\circ}\text{C)}$ and $R = 1.987 \text{ cal} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$.

Further *trans*-acting *delta* ribozyme variants were produced using plasmid p δ RzP1.1. The variants have either A23 or C24 mutated to one of the other three possible bases. The six resulting *delta* ribozyme variants are named for the altered nucleotide (δ RzP1-A23C, -A23G, -A23U, -C24A, -C24G, and -C24U; Table 3). Complementary or compensatory substrates (Table 3) were generated in which either position 7 or 8 of the wild type substrate (SP1.1) was altered in order to restore the Watson-Crick base pair formation of the P1 stem between the substrates and the ribozyme variants.

TABLE 3

Transcripts	Sequence
Substrates	
SP1.1	₁ GGGCGGGUCGG ₁₁
SG7A	GGGCGGA <u>A</u> UCGG
SG7C	GGGCGGC <u>C</u> UCGG
SG7U	GGGCGGU <u>U</u> UCGG
SU8A	GGGCGGG <u>A</u> CGG
SU8C	GGGCGGG <u>C</u> CGG
SU8G	GGGCGGG <u>G</u> CGG
SU8G-9mers	₁ GCGGGG <u>G</u> CGG ₉
Ribozymes	
δ RzP1.1	₂₀ CCGACCU ₂₆
δ RzP1-A23C	CCG <u>C</u> CCU
δ RzP1-A23G	CCG <u>G</u> CCU
δ RzP1-A23U	CCG <u>U</u> CCU
δ RzP1-C24A	CCGA <u>A</u> CU
δ RzP1-C24G	CCGAG <u>C</u> U
δ RzP1-C24U	CCGAU <u>C</u> U

The extent of cleavage of the δ RzP1-C24N ribozyme variants were compared with that of the wild type ribozyme δ RzP1.1 for each of 4 substrates (A), and correspondingly, the extent of cleavage of the δ RzP1-C24N ribozyme variants were

Complementary pairs of substrates and ribozymes were used for kinetic studies to obtain the experimental data required for the calculation of apparent K_m (K_m') and apparent k_2 values and the results are shown in Table 4.

06-000000

A collection of 13 substrates including all single mutants for positions -4 to -1 compared to the original substrate were synthesized. Positions -4 to -1 refer to the four nucleotides directly 5' to the cleavage site, position -1 being right next to the cleavage site and position -4 being the furthest from the cleavage site, as shown in Figure 2. For each mutant, trace amounts of 5'-³²P-labeled substrates (<1 nM) were incubated in the presence of an excess of ribozyme (200 nM), and the maximal cleavage percentages (i.e. end-point) (pre-steady state conditions) determined as a comparative parameter. The Applicant observed that the base requirement varies for each position. At position -1, the base preference was A > C > U >> G, where a guanosine at this position rendered the substrate uncleavable. At position -2, an A improved the cleavage efficiency compared to the original G, while a substrate with a U was poorly cleaved and a C gave an uncleavable substrate. In contrast at position -3, C, U and A gave substrates that have a two fold improved cleavage compared to the wildtype G. Finally at position -4, the presence of a pyrimidine (i.e. C or U) improved the maximal percentage of cleavage by at least two fold compared to a purine (i.e. G or A).

25 In order to assess accurately the base requirement
at each position, kinetic analysis were performed under pre-
steady-state conditions. Pseudo first-order cleavage rate
constants (k_2 and K_m') were measured with an excess of ribozyme
(5 to 600 nM) and trace amounts of end-labeled substrate (<0.1
30 nM).

TABLE 5

Position	Identity	K_m' (nM)	k_2 (min ⁻¹)	k_2/K_m' (nM ⁻¹ min ⁻¹)	Specificity index
-1	C	31.52	0.22	6.66×10^{-3}	1.00
	U	33.2	0.11	3.34×10^{-3}	0.50
	A	14.27	0.27	1.79×10^{-2}	2.68
	C	na	na	na	na
-2	G	31.52	0.22	6.66×10^{-3}	1.00
	A	28.7	0.33	1.15×10^{-4}	1.73
	C	na	na	na	na
	U	94	0.08	8.19×10^{-4}	0.12
-3	C	31.52	0.22	6.66×10^{-3}	1.00
	A	9.93	0.20	1.99×10^{-2}	3.02
	C	11.3	0.24	2.10×10^{-2}	3.15
	U	8.76	0.20	2.32×10^{-2}	3.48
-4	G	31.52	0.22	6.66×10^{-3}	1.00
	A	27.14	0.12	4.45×10^{-3}	0.67
	C	11.81	0.27	1.86×10^{-2}	2.79
	U	16.42	0.23	1.40×10^{-2}	2.10

Table 5. Kinetic analysis of the collection of single mutated substrates. Pseudo first-order cleavage rate constants (k_2 and K_m') were measured using an excess of ribozyme (5 to 600 nM) and trace amounts of end-labelled substrate (< 0.1 nM). Apparent second-order rate constants (k_2/K_m') were calculated and their relative specificity determined as compared to the original substrate. The values were calculated from at least two independent experiments, and errors were less than 25%. Sequence for position -4 to -1 are indicated for each substrate.

Then, apparent second-order rate constants (k_2/K_m') were calculated and a specificity index determined, fixing arbitrarily as 1.00 the values of the original substrate (i.e. $\text{.}_4\text{GGGC}_1$). At position -1, the presence of a uridine resulted in a similar relative specificity (0.50) while the presence of an adenine increased the relative specificity to 2.68. This increase appears mainly as a result of a K_m' decrease of 2 fold. For position -2, the presence of a purine (i.e. G or A)

gave similar relative specificity (1.73, compared to 1.00, respectively). In contrast, the presence of a uridine resulted in a poorly cleaved substrate, while when a cytosine was present, the substrate was uncleavable. In the case of the uridine at position -2, the specificity was evaluated to be reduced from 8 fold to 0.12 compared to the original substrate (i.e. 1.00). The decrease in specificity appears to result from a 3 fold increase of the K_m' and a 3 fold decrease of the k_2 value. These results show a clear preference for purine in position -2, and a pyrimidine should be avoided in that position.

For position -3, when the guanosine of the original substrate was replaced by any other base (i.e. A, C, or U), the K_m' was lowered by 3 fold while the k_2 remained almost identical, resulting in an specificity increase ranging from 3.02 to 3.48. Finally for position -4, a purine (G and A) yield a substrate with about the same specificity (i.e. 0.67 and 1.00). However, the presence of a pyrimidine in position -4 improved the specificity by at least two fold with 2.79 and 2.10 for a C and a U, respectively. Specifically, the presence of a C or a U the K_m' was lowered, while the k_2 remained almost identical. Thus, it appears clear that the base requirement from position -4 to -1 of the substrate, contributes significantly and differently to the ability of the substrate to be cleaved.

Based on the observation that mutations in position -3 were those that most strongly increased the relative specificity, the Applicant investigated whether or not a larger amount of Mg^{2+} in the cleavage reaction would affect the kinetic parameters of these substrates. Under single turnover conditions, in which the ribozyme and substrate concentrations were kept at 200 nM and 1 nM, respectively, the Applicant found that the ribozyme cleaved these substrates at Mg^{2+} concentrations as low as 1 mM, which is the estimated physiological concentration of Mg^{2+} (Ananovorovich, S. and Perreault, J.P. (1998) *J. Biol. Chem.*, **273**, 13182-13188, and Trut, T.W. (1994) *Mol. Cell. Biochem.*, **140**, 1-22). A maximum

k_{obs} for each substrate was observed when the concentration of Mg^{2+} was 10 mM. The requirement for magnesium at half-maximal velocity (K_M) was similar for these mutated substrates and the original substrate, varying between 1.5 to 2.2 mM. Similar experiments were also performed with several other substrates from the collection and identical results were obtained, suggesting that the differences of the kinetic parameters for various substrates were not related to different affinity for the magnesium.

Notably, the cleavage assays performed with the initial collection of substrates (i.e. single mutants) indicated that the presence of a pyrimidine in the position -2 either reduces the cleavage activity or yields an uncleavable substrate. Specifically, a uridine decreases the relative specificity by 8 fold while a cytosine inhibits the cleavage completely (see Table 6). One plausible explanation of such results is that when a C is present at position -1 and followed by a pyrimidine (i.e. C or U) at position -2, both nucleotides of the substrate may interact with nucleotides located on the ribozyme resulting in inactive substrate/ribozyme complex. It seems reasonable to suggest that base-pairing may be formed with the ribozyme guanosines at position 27 and 28 of the J1/4 junction, which new base pairs will compete with formation of the P1.1 stem (Fig. 2). In this case, a cytosine in position -2 will form two consecutive GC base pairs. Similarly, a uridine in position -2 allows formation of a GC follow by a GU, which will be less stable than two GC's, yielding a reduced activity compared to the absence of activity. In order to learn more about the nucleotide preference in position -2, taking into account the neighboring positions, a second collection of substrates with more than one mutation were synthesized.

First, the Applicant verified whether a cytosine at position -2 after non-cytosine at position -1 has a detrimental effect. Based on the previous results, a substrate with an adenine in position -1 and a cytosine in position -2, S-A₁C₂, was synthesized and further tested for cleavage

0969967-103000

efficiency. A moderate extent of cleavage of 14% was observed at 200 nM ribozyme, which is less than the substrates including either the sequence C₁G₂ or A₁G₂. In comparison to the substrate with the sequence A₁G₂, the S-A₁C₂ substrate showed a virtually identical apparent K_m (K_m') while the cleavage constant (k₂) was reduced by approximately 4 fold, yielding a 4-fold reduction of the relative specificity (i.e. from 2.68 to 0.60; Table 6). These results suggest that the presence of a cytosine at position -2 reduces significantly the cleavage of a substrate. Moreover, if this cytosine is followed by a second cytosine in position -1, the result is an uncleavable RNA molecule (see above).

TABLE 6

Mutant	K _m ' (nM)	k ₂ (min ⁻¹)	K ₂ /K _m ' (nM ⁻¹ min ⁻¹)	Specificity index
SC ₁ G ₂	31.5	0.22	6.98 x 10 ⁻³	1
SA ₁	14.3	0.27	1.89 x 10 ⁻²	2.68
SA ₁ C ₂	15.4	0.06	3.9 x 10 ⁻³	0.6
SA ₁ C ₂ C ₃	15.2	0.039	2.57 x 10 ⁻³	0.4
SA ₁ C ₂ C ₃ C ₄	16.5	0.25	1.52 x 10 ⁻²	2.28

Table 6. Kinetic analysis of the collection of multiple mutated substrates. Pseudo first-order cleavage rate constants (k₂ and K_m') were measured using an excess of ribozyme (5 to 600 nM) and trace amounts of end-labelled substrate (<0.1 nM). Apparent second-order rate constants (k₂/K_m') were calculated and their relative specificity determined as compared to the original experiments, and errors were less than 25%. Sequence for position -4 to -1 are indicated for each substrate.

Secondly, the Applicant verified whether a cytosine at position -2 followed by a cytosine at position -3 gives a cleavable substrate. In other words, two consecutive cytosines, regardless of their positions, will yield uncleavable substrates. Therefore, the Applicant synthesized the substrate S-A₁C₂C₃ and verified its ability to be cleaved. The S-A₁C₂C₃ put together was cleaved with kinetic parameters

almost identical to the the substrate S-A₁C₂ substrate except that the k_2 was slightly reduced to 0.039 min⁻¹ compared to 0.062 min⁻¹, resulting in a small reduction of the relative specificity (i.e. from 0.60 to 0.40; Table 6). These results show that the presence of a cytosine at position -3 following a cytosine at position -2 reduced slightly the cleavage activity, and did not significantly modify the ability of a substrate to be cleaved. Thus, a cytosine at position -3 does not have the same influence as that at position -2.

Thirdly, the Applicant asked whether two consecutive cytosines at positions -4 and -3 give a similar effect yielding uncleavable (or less cleaved) substrate. A substrate containing cytosines at positions -3 and -4 and adenines in position -1 and -2 was synthesized. Adenines were included in position -1 and -2 because this residue appears to give a readily cleaved substrate as compared to the single mutation collection (see above). The S-A₁A₂C₃C₄ mutant has a maximum cleavage of 61%. Moreover, the Applicant determined a K_m' of 16.5 nM and a k_2 value increased to 0.25 min⁻¹, resulting in a substrate with a relative specificity of 2.28 as compared to the original substrate (Table 6). Thus, the presence of two consecutive cytosines at position -3 and -4 has no detrimental effect.

Finally, the Applicant asked whether it is possible to compensate for the detrimental effect of the presence of two consecutive cytosines at positions -1 and -2, by including the one at position -2 in a hairpin structure. A longer RNA substrate (i.e. 18-mer compared to 14mer) including a hairpin at 5'-end, which involved the C₂ in the last base pair of the helix was chemically synthesized and then tested. This substrate was poorly cleaved. Only trace amounts of products were detected (i.e. maximum percentage cleavage of <2.0 %), and as a consequence, no more extensive characterization was possible. If the sequence was drawn in order to avoid the formation of the 5'-end hairpin (i.e. C₂ remains single strand; S-hp-), no cleavage at all was observed. These two results showed that the presence of a base-paired cytosine at

position -2 gave minimal activity as compared to this cytosine in single strand. However, the improvement was very limited.

Example 4: L4 loop Modifications

5 A modified form of δ RzP1.1 described above was made by replacing the L4 loop sequence GCUU which is relatively unstable, with the ultrastable L4 loop (UUCG) (shown on the right in Figure 3). The kinetic parameters (k_{cat} and K_M) and dissociation constant (K_d) were virtually identical.

10

Example 5: Bimolecular ribozyme

A modified form of δ RzP1.1 described above was made by dividing the L4 loop into two resulting in two fragments, namely, RzA and RzB (as shown in Figure 4). The RzA consists
15 of 37 nucleotides encompassing a substrate recognition site (P1 stem), P3 stem and portions of P2 and P4 stems. The RzB consists of 20 nucleotides which is able to base pair to RzA to form a bimolecular ribozyme complex. RzA and RzB were synthesized as described in Example 1. Because both RzA and
20 RzB are relatively small, they can be chemically synthesized. Therefore, this bimolecular delta ribozyme allows the introduction of any chemically modified nucleoside.

Example 6: Deoxyribonucleotide modifications

25 Example 5 describes a bimolecular ribozyme. Modified versions of the ribozyme described in Example 5 were made by replacing one ribonucleotide in RzB with a deoxyribonucleotide individually at positions 9 to 15. This resulted in 7 different RzB's each containing one
30 deoxyribonucleic acid.

Sub (12)
35 The influence of 2'-OH groups in RzB on the catalytic activity of RzA:RzB complex was analyzed. 0.066 μ M of a mix of cold and end-labeled RNA substrates were incubated in presence of 0.066 μ M of RzA and 0.2 μ M of various RzB RNA/DNA mixed polymers. The incubation was performed in 50 mM Tris-HCl pH 8.0 and 50 mM $MgCl_2$ at 37°C. An aliquot was

removed after one hour and the reaction stopped by the addition of an excess of stop solution (xc, bb, formamide). Reaction mixtures were fractionated on 20% polyacrylamide gel electrophoresis and were exposed on x-ray films. Fully deoxyribonucleotide RzB molecules are not able to support a cleavage activity. Individual deoxy substitution mutants were subjected to catalytic cleavage. All of the reconstituted complexes were active to different extents. S and P respectively represent substrate and product species. As an example, dGg stands for GGCgCAUGgCUAAGGGACCC where uppercase and lowercase letters respectively represent ribo- and deoxyribonucleotides. The results are shown in Figure 6 and Table 7.

Table 7 shows the quantification of time course experiments performed. Rate and extent of cleavage values were obtained from fitting the experimental data to the equation $A_t = A_{\alpha}(1 - e^{-kt})$ where A_t is the percentage of cleavage at time, t , A_{α} is the maximum cleavage and k is the reaction rate. Data analysis was performed with GraFit Version 3.01 from Erithacus Software.

TABLE 7

Species	Rate (min^{-1})	Extent (%)
RzB	5.7×10^{-2}	27.01
dG9	3.3×10^{-2}	9.80
dC10	2.4×10^{-2}	30.42
dU11	4.6×10^{-2}	45.87
dA12	4.0×10^{-2}	26.79
dA13	1.8×10^{-2}	27.46
dG14	8.0×10^{-2}	61.44
dG15	7.8×10^{-2}	54.15

Table 7. Rate and extent of substrate cleavage using 2'-OH modified ribozymes.

5

~~Plasmids encoding the HDag mRNA and delta ribozymes.~~

35 *RNA Synthesis.* *In vitro* transcription: HDAG mRNA
was transcribed from *Hind* III-linearized pKSagS, while
ribozymes were transcribed from *Sma* I-linearized ribozyme

0996710000

[illegible]

Oligonucleotide probing. DNA oligonucleotides complementary to the potential target sites were purchased from Gibco-BRL and 5'-end labelled using T4 polynucleotide kinase (Pharmacia) in the presence of 10 μ Ci [γ - 32 P]ATP. Labelled oligonucleotides (~ 2 500 cpm; ~ 0.05 nM) and unlabelled mRNA (2.4 to 1 200 nM) were hybridized together for 10 min at 25°C in a solution containing 50 mM Tris-HCl pH 7.5 and 10 mM MgCl₂ in a final volume of 15 μ l. Loading solution (2 μ l of 1X TBE, 10 mM MgCl₂, 40% glycerol, 0.25% bromophenol blue and 0.25% xylene cyanol) was added, and the resulting solutions fractionated on native 5% PAGE gels (30:1 ratio of acrylamide to bisacrylamide, 50 mM Tris-borate pH 8.3, 10 mM MgCl₂ and 5% glycerol) at 4°C in the presence of recirculating 50 mM Tris-borate pH 8.3 and 10 mM MgCl₂ buffer. The dried gels were analyzed with the aid of a PhosphorImager (Molecular Dynamics). RNase H probing was performed using the same oligonucleotides. In these experiments randomly labelled S-HDAg mRNA (~10 000 cpm; ~10 nM) and unlabelled oligonucleotides (1 μ M) were annealed as described for gel shift assays for 10 min, then 0.2 U of *E. coli* RNase H (Pharmacia) was added and the reaction incubated at 37°C for 20 min. The reactions were stopped by the addition of stop-solution (3 μ l of 97% formamide, 10 mM EDTA, 0.25% bromophenol blue and 0.25% xylene cyanol), fractionated on 5% denaturing PAGE gels, and analyzed by autoradiography.

In vitro cleavage assays and kinetic analyses.

Cleavage assays were performed at 37°C under single turnover conditions with either randomly labelled mRNA (~ 10 nM) or 5'-end labelled small substrates (<1nM), and an excess of ribozyme (2,5 µM) in 10 µL final volume containing 50 mM Tris-HCl pH 8.0 and 10 mM MgCl₂. A pre-incubation of 5 min at 37°C preceded the addition of the Tris-magnesium buffer which initiates the reaction. After an incubation of 1 to 3 hrs at 37°C, stop-solution (5µl) was added and the mixture quickly

30
SUB
CS

stored at -20°C until its fractionation on 5% denaturing PAGE gels and subsequent autoradiography. Cleavage sites of the active ribozymes were verified by primer extension assays as described previously (Côté, F. and Perreault, J.-P. (1997) *J. Mol. Biol.*, **273**, 533-543). Briefly, oligonucleotides were synthesized to have complementary sequence to positions downstream (~ 100 positions) from the cleavage site according to the mRNA. For example, for the cleavage site of Rz-12, the oligonucleotide primer, 5'CTTTGATGTTCCCCAGCCAGG-3' (21mer),
10 was used in the reverse transcriptase reaction containing the ribozyme cleavage reaction mixture.

Active ribozymes (Rz-1, -11 and -12) were characterized under single turnover conditions essentially as described in Example 1.

0969667-103000

5
C
A
S

TABLE 8

Ribozyme	P1 stem sequence	Size of expected cleavage products (nt)
Rz-1	CCCAGCU	265 , 551
Rz-2	CCUCUUU	330 , 486
Rz-3	CCUUGUU	403 , 413
Rz-4	UGUUCUU	440 , 376
Rz-6	GGGGUUU	572 , 244
Rz-7	UCCCCUU	593 , 223
Rz-9	GGACUCU	640 , 176
Rz-11	UCGACUU	130 , 686
Rz-12	GCCACCU	175 , 641

Sub
Ne TABLE 8 (cont'd)

mRNA sequence

1	CACCGCGGU	GCGGCGC	UAGAACUAG	GGAUCCCU	GGCUCGGGCG	GCGAGUCC
61	CAGUCUCCU	UUUACAGA	AUGUAAGAG	ACUGAGGA	GCCGCCUCUA	GCCGAGAU
121	GCCGGUCCG	<u>GUCGAGGA</u>	AACCGCGGA	GGAGAGAA	GAUCCUCGAG	CAGUGGGU (11, 12)
181	<u>CCGGAAGAA</u>	GAAGUUAG	GAACUCGAG	GAGACCUC	GAAGACAAAG	AAGAAACU
241	AGAAGAUAG	GGACGAAA	CCCUGGCUG	GGAACAUC	AGGAAUUCUC	GGAAAGAA (1)
301	AUAAGGAUG	AGAGGGG	CCCCCCGCG	AGAGGGCC	AACGGACCAG	AUGGAGGU (2)
361	ACUCCGGAC	UCGGAAGA	CCUCUCAGG	GAGGAUUC	<u>CGACAAGGAG</u>	AGGCAGGA (3)
421	CCGACGAAG	AAGGCCCU	<u>AGAAACAAGA</u>	GAAGCAGC	UCGGCGGGAG	GCAAGAAC (4)
481	CAGCAAGGA	GAAGAAGA	AACUCAGGA	GUUGACCG	GAAGACGAGA	GAAGGGAA
541	AAGAGUAGC	GGCCCGCC	UUGGGGGUG	<u>GAACCCCC</u>	GAAGGUGGAU	<u>CGAGGGGA</u> (6, 7)
601	GCCCGGGGG	GGCUUCGU	CCAAUCUGC	GGGAGUCC	<u>GAGUCCCCCU</u>	UCUCUCGG (9)
661	CGGGGAGGG	CUGGACAU	GGGGAACC	GGGAUUUC	UAGGAUAUAC	UCUUCCCA
721	CGAUCCGCC	UUUUCUCC	AGAGUUUGC	ACCCAGU	AUAAAGCGGG	UUUCCACU
781	CAGGUUUGC	UCUGCGGU	UUCUUUCCU	UUC		

Table 8. Synthesized delta ribozyme. Previous page is the ribozyme nomenclature with the sequence composing the P1 stem domain and the size of the expected products. This page is the mRNA sequence. The mRNA sequences targeted by ribozymes are underlined, and the ribozyme number is in parentheses on the right.

Example 8: Cleavage Assay of a ribozyme of the invention
10 against 552 nt-HBV RNA substrate.

500 nM of a delta ribozyme as shown in Figure 7 was incubated with 1 nM randomly-labelled 552 nt-HBV (human hepatitis B virus) mRNA at 37°C in the presence of 50 mM Tris-HCl pH 7.5 and 10 mM MgCl₂. A single exponential equation was used to fit data to $k_{\text{obs}} = 0.031 \text{ min}^{-1}$ with 28% cleavage. This demonstrates that a ribozyme of the invention cleaves mRNA from the human hepatitis B virus.

1. **Introduction**
 2. **Background**
 3. **Methodology**
 4. **Results**
 5. **Discussion**
 6. **Conclusion**
 7. **References**
 8. **Appendix**
 9. **Figure 1**
 10. **Figure 2**
 11. **Figure 3**
 12. **Figure 4**
 13. **Figure 5**
 14. **Figure 6**
 15. **Figure 7**
 16. **Figure 8**
 17. **Figure 9**
 18. **Figure 10**
 19. **Figure 11**
 20. **Figure 12**
 21. **Figure 13**
 22. **Figure 14**
 23. **Figure 15**
 24. **Figure 16**
 25. **Figure 17**
 26. **Figure 18**
 27. **Figure 19**
 28. **Figure 20**
 29. **Figure 21**
 30. **Figure 22**
 31. **Figure 23**
 32. **Figure 24**
 33. **Figure 25**
 34. **Figure 26**
 35. **Figure 27**
 36. **Figure 28**
 37. **Figure 29**
 38. **Figure 30**
 39. **Figure 31**
 40. **Figure 32**
 41. **Figure 33**
 42. **Figure 34**
 43. **Figure 35**
 44. **Figure 36**
 45. **Figure 37**
 46. **Figure 38**
 47. **Figure 39**
 48. **Figure 40**
 49. **Figure 41**
 50. **Figure 42**
 51. **Figure 43**
 52. **Figure 44**
 53. **Figure 45**
 54. **Figure 46**
 55. **Figure 47**
 56. **Figure 48**
 57. **Figure 49**
 58. **Figure 50**
 59. **Figure 51**
 60. **Figure 52**
 61. **Figure 53**
 62. **Figure 54**
 63. **Figure 55**
 64. **Figure 56**
 65. **Figure 57**
 66. **Figure 58**
 67. **Figure 59**
 68. **Figure 60**
 69. **Figure 61**
 70. **Figure 62**
 71. **Figure 63**
 72. **Figure 64**
 73. **Figure 65**
 74. **Figure 66**
 75. **Figure 67**
 76. **Figure 68**
 77. **Figure 69**
 78. **Figure 70**
 79. **Figure 71**
 80. **Figure 72**
 81. **Figure 73**
 82. **Figure 74**
 83. **Figure 75**
 84. **Figure 76**
 85. **Figure 77**
 86. **Figure 78**
 87. **Figure 79**
 88. **Figure 80**
 89. **Figure 81**
 90. **Figure 82**
 91. **Figure 83**
 92. **Figure 84**
 93. **Figure 85**
 94. **Figure 86**
 95. **Figure 87**
 96. **Figure 88**
 97. **Figure 89**
 98. **Figure 90**
 99. **Figure 91**
 100. **Figure 92**
 101. **Figure 93**
 102. **Figure 94**
 103. **Figure 95**
 104. **Figure 96**
 105. **Figure 97**
 106. **Figure 98**
 107. **Figure 99**
 108. **Figure 100**
 109. **Figure 101**
 110. **Figure 102**
 111. **Figure 103**
 112. **Figure 104**
 113. **Figure 105**
 114. **Figure 106**
 115. **Figure 107**
 116. **Figure 108**
 117. **Figure 109**
 118. **Figure 110**
 119. **Figure 111**
 120. **Figure 112**
 121. **Figure 113**
 122. **Figure 114**
 123. **Figure 115**
 124. **Figure 116**
 125. **Figure 117**
 126. **Figure 118**
 127. **Figure 119**
 128. **Figure 120**
 129. **Figure 121**
 130. **Figure 122**
 131. **Figure 123**
 132. **Figure 124**
 133. **Figure 125**
 134. **Figure 126**
 135. **Figure 127**
 136. **Figure 128**
 137. **Figure 129**
 138. **Figure 130**
 139. **Figure 131**
 140. **Figure 132**
 141. **Figure 133**
 142. **Figure 134**
 143. **Figure 135**
 144. **Figure 136**
 145. **Figure 137**
 146. **Figure 138**
 147. **Figure 139**
 148. **Figure 140**
 149. **Figure 141**
 150. **Figure 142**
 151. **Figure 143**
 152. **Figure 144**
 153. **Figure 145**
 154. **Figure 146**
 155. **Figure 147**
 156. **Figure 148**
 157. **Figure 149**
 158. **Figure 150**
 159. **Figure 151**
 160. **Figure 152**
 161. **Figure 153**
 162. **Figure 154**
 163. **Figure 155**
 164. **Figure 156**
 165. **Figure 157**
 166. **Figure 158**
 167. **Figure 159**
 168. **Figure 160**
 169. **Figure 161**
 170. **Figure 162**
 171. **Figure 163**
 172. **Figure 164**
 173. **Figure 165**
 174. **Figure 166**
 175. **Figure 167**
 176. **Figure 168**
 177. **Figure 169**
 178. **Figure 170**
 179. **Figure 171**
 180. **Figure 172**
 181. **Figure 173**
 182. **Figure 174**
 183. **Figure 175**
 184. **Figure 176**
 185. **Figure 177**
 186. **Figure 178**
 187. **Figure 179**
 188. **Figure 180**
 189. **Figure 181**
 190. **Figure 182**
 191. **Figure 183**
 192. **Figure 184**
 193. **Figure 185**
 194. **Figure 186**
 195. **Figure 187**
 196. **Figure 188**
 197. **Figure 189**
 198. **Figure 190**
 199. **Figure 191**
 200. **Figure 192**
 201. **Figure 193**
 202. **Figure 194**
 203. **Figure 195**
 204. **Figure 196**
 205. **Figure 197**
 206. **Figure 198**
 207. **Figure 199**
 208. **Figure 200**
 209. **Figure 201**
 210. **Figure 202**
 211. **Figure 203**
 212. **Figure 204**
 213. **Figure 205**
 214. **Figure 206**
 215. **Figure 207**
 216. **Figure 208**
 217. **Figure 209**